

# Willowcreek Animal Hospital Vaccination Program

Livestock Type-Date--	2-3 weeks prior to calving	Calving	Branding	3-4 weeks prior to breeding	2-3 weeks prior to weaning or weaning	Weaning Or 2-3 weeks post weaning	Preg-checking
<b>CALVES-</b>		Dip Navel BoSe 1mL	Vista 5 Vision8 OncePMH IN +/- Implant Ralgrow		Vista Once  Vision8  Deworm	Vista5  Vision8	
<b>Replacement HEIFERS-</b>	Guardian  Booster 3-4 weeks			Vista5VL5 Vision8 Booster Vista5VL5 3 weeks later  +/- Deworm			Vision8  +/- Deworm/ Delice
<b>COWS-</b>	Guardian			Vista5VL5  +/-Deworm			Vision8  +/- Deworm/ Delice
<b>BULLS-</b>		Breeding Soundness Exam  Trich	Vista5VL5  Vision8  +/- Deworm				+/-Deworm/ Delice
<b>Guidelines-</b>	Read and follow label directions.	Place shots in neck when possible.	Never combine vaccines. Avoid off-label or extra label use.	Keep vaccines out of sunlight and in a cool place.	Choose the correct needle. Use transfer needles.	Don't mix too much. Shake larger dosage bottles to ensure consistent dose.	Do not use disinfectants with modified live product.

*Willowcreek Animal Hospital (208) 529-4081 Fremont Animal Care Center (208) 624-7218*